

Taxol in the Management of Ovarian Cancer

Jung-Min Lee, MD

NIH Lasker Clinical Research Scholar
Women's Malignancies Branch, CCR, NCI

Disclosures

I have no remunerated positions.


**I may answer questions with responses indicating
investigational use of agents or use of
investigational agents.**

Rosalind Franklin

- Received her PhD from Cambridge in 1945
- Early 1950s, discovery of DNA structure
- 1956, diagnosed with ovarian cancer
- 1958, died in London




(a) Rosalind Franklin




(b) Franklin's X-ray diffraction photograph of DNA

Ovarian cancer survival trends




Have our interventions been successful?

US Cancer Death Rates 1930-2008



Slowly but surely, we have made a good progress



Ovarian cancer

Rare but most lethal gynecologic cancer in the U.S.

Estimated New Cases

		Males	Females		
Prostate	161,360	19%		Breast	252,710 30%
Lung & bronchus	116,990	14%		Lung & bronchus	105,510 12%
Colon & rectum	71,420	9%		Colon & rectum	64,010 8%
Urinary bladder	60,490	7%		Uterine corpus	61,380 7%
Melanoma of the skin	52,170	6%		Thyroid	42,470 5%
Kidney & renal pelvis	40,610	5%		Melanoma of the skin	34,940 4%
Non-Hodgkin lymphoma	40,080	5%		Non-Hodgkin lymphoma	32,160 4%
Leukemia	36,290	4%		Leukemia	25,840 3%
Oral cavity & pharynx	35,720	4%		Pancreas	25,700 3%
Liver & intrahepatic bile duct	29,200	3%		Kidney & renal pelvis	23,380 3%
All Sites	836,150	100%		All Sites	852,630 100%

**Estimated
22,400 new cases
(2.6%) in 2017**



Estimated Deaths

		Males	Females		
Lung & bronchus	84,590	27%		Lung & bronchus	71,280 25%
Colon & rectum	27,150	9%		Breast	40,610 14%
Prostate	26,730	8%		Colon & rectum	23,110 8%
Pancreas	22,300	7%		Pancreas	20,790 7%
Liver & intrahepatic bile duct	19,610	6%		Ovary	14,080 5%
Leukemia	14,300	4%		Uterine corpus	10,920 4%
Esophagus	12,720	4%		Leukemia	10,200 4%
Urinary bladder	12,240	4%		Liver & intrahepatic bile duct	9,310 3%
Non-Hodgkin lymphoma	11,450	4%		Non-Hodgkin lymphoma	8,690 3%
Brain & other nervous system	9,620	3%		Brain & other nervous system	7,080 3%
All Sites	318,420	100%		All Sites	282,500 100%

**Fractional death
rate 63%**



“Ovarian” cancer histology

High grade serous




Low grade serous

Mucinous




Endometrioid

Clear cell



Genetic sources of genomic instability

Mutually exclusive events



**Treatment with Taxol still matters for ovarian cancer
in this genomic era**

Introduction of Taxol to the treatment of ovarian cancer

[CANCER RESEARCH 47, 2486–2493, May 1, 1987]

Phase I Clinical and Pharmacokinetic Study of Taxol¹

Peter H. Wiernik,² Edward L. Schwartz, Janice J. Strauman, Janice P. Dutcher, Richard B. Lipton, and Elisabeth Paietta

Albert Einstein Cancer Center, Montefiore Medical Center, Bronx, New York 10467

Introduction of Taxol to the treatment of ovarian cancer

[CANCER RESEARCH 47, 2486–2493, May 1, 1987]

Phase I Clinical and Pharmacokinetic Study of Taxol¹

Peter H. Wiernik,² Edward L. Schwartz, Janice J. Strauman, Janice P. Dutcher, Richard B. Lipton, and Elisabeth Paietta

Albert Einstein Cancer Center, Montefiore Medical Center, Bronx, New York 10467

ARTICLES

Taxol: A Unique Antineoplastic Agent with Significant Activity in Advanced Ovarian Epithelial Neoplasms

William P. McGuire, MD; Eric K. Rowinsky, MD; Neil B. Rosenshein, MD;
Francis C. Grumbine, MD; David S. Ettinger, MD; Deborah K. Armstrong, MD;
and Ross C. Donehower, MD

FDA approval of taxol for ovarian cancer in 1992


30% response rate in platinum-resistant ovarian cancer
Annals of Internal Medicine. 1989;111:273-279.

Changing the treatment paradigm

Taxol plus platinum agent is superior to other combo

GOG111: Ovarian (suboptimal III/IV)


- Epithelial Ovarian Cancer
- Suboptimal Stage III/IV
- No prior therapy




Open: 13-Apr-90
Closed: 02-Mar-92
Accrual: 410 pts

McGuire, et al. NEJM 334:1-6, 1996

GOG 182: less is better



It does not matter whether you give before or after surgery




Vergote I et al, NEJM, 2010

CONFIRMED: Kehoe S et al, Lancet Oncol 2015

Pooled analysis EORTC55971 and CHORUS Vergote et al. Lancet Oncol 2018

JGOG: dose-dense weekly Paclitaxel

- Epithelial ovarian or peritoneal (including clear cell/mucinous)
- Stage II – IV
- Stratified: residual disease, stage, histology
- Primary endpoint: PFS
- Secondary: OS



- Dose-dense paclitaxel hematologic toxicity, fewer pts completed protocol therapy
- Improved PFS and OS in Japanese patients with dose-dense weekly paclitaxel


Accrual: 637 pts (intent-to-treat)

Katsumata N, et al Lancet, 2009
Katsumata N, et al Lancet Oncol, 2013

GOG172: IP vs. IV

The way of delivery of carboplatin and taxol may matter

- EOC
- Optimal stage III
- Elective SLO




Upfront treatment summary

- **Surgery followed by taxane/platinum-based chemotherapy is standard treatment for newly diagnosed ovarian cancer patients**
- **Taxol plus platinum-based chemotherapy combination is active in ovarian cancer, independent of histology**
- **IP chemotherapy would benefit the selected population**

Taking to the next level

PARP inhibitor maintenance following chemotherapy



**SOLO1: olaparib maintenance
in BRCA mutant patients
FDA approval 2018**

	Olaparib (n=260)	Placebo (n=131)
Events (%)[50.6% maturity]	102 (39.2)	96 (73.3)
Median PFS, months	NR	13.8
	HR 0.30	



95% CI 0.23 to 0.41 P<0.0001

Moore et al. NEJM 2018

Taking to the next level

Bevacizumab maintenance following chemotherapy

- EOC
- III/IV




GOG218: bevacizumab maintenance
FDA approval 2018

FDA approval June 2018


Unfortunately, recurrence is common in ovarian cancer



AURELIA: Chemo + bevacizumab vs chemo Progression-free survival




AURELIA: weekly paclitaxel drove results



Chemo arm*	ORR (bev/no bev)	PFS HR (bev/no bev)	OS HR (bev/no bev)
Paclitaxel	53.3 v 30.2%	0.46 (10.4 v 3.9 mo)	0.65 (22.4 v 13.2 mo)
PLD	17.0 v 0%	0.57 (5.4 v 3.5 mo)	0.91 (13.7/14.1 mo)
topotecan	13.7 v 7.8%	0.32 (5.8 v 2.1 mo)	1.09 (13.8/13.3 mo)


Making another history

- Immunomodulatory effect of Taxol



Making another history

- Immunomodulatory effect of Taxol




- Ongoing phase 3 clinical trials in the upfront treatment setting using Taxol/Carboplatin + immune checkpoint inhibitor +/- anti-angiogenic drug

Other new directions in the treatment of ovarian cancer

Targeted agents combination


PARP inhibitor olaparib and VEGF receptor inhibitor cediranib in Phase III Registration Trials

**Platinum-sensitive
Ovarian cancer
NRG GY-004**



**Platinum-refractory/resistant
Ovarian Cancer
NRG GY-005 (PI: J Lee)**


Anticipated completion of accrual: Sep 2020



* Weekly Taxol, topotecan or PLD

First-in-human combination of PARP inhibition + anti-PD-L1

Phase I olaparib + durvalumab in recurrent ovarian cancer



Conclusions

- Ovarian cancer is one of the leading cause of death in the industrialized countries
- The standard of care includes surgical resection and paclitaxel-platinum based chemotherapy
- IP chemotherapy would benefit the selected population

Conclusions

- Ovarian cancer is one of the leading cause of death in the industrialized countries
- The standard of care includes surgical resection and paclitaxel-platinum based chemotherapy
- IP chemotherapy would benefit the selected population
- Ovarian cancer recurrence is common
- Paclitaxel is active either in combination with bevacizumab or as monotherapy in recurrent setting
- There are still unmet needs for highly effective less toxic therapies



NATIONAL CANCER INSTITUTE
Center for Cancer Research

ccr.cancer.gov